

Impact of the second order self-forces on the dephasing of the gravitational waves from quasi-circular extreme mass-ratio inspirals

Soichiro Ioyama^{1,*}, Ryuichi Fujita^{2,†}, Norichika Sago^{3,‡}, Hideyuki Tagoshi^{4,§} and Takahiro Tanaka^{1,¶}

¹ Yukawa Institute for Theoretical Physics, Kyoto university, Kyoto, 606-8502, Japan

² Departament de Física, Universitat de les Illes Balears, Palma de Mallorca, E-07122 Spain

³ Faculty of Arts and Science, Kyushu University, Fukuoka 819-0395, Japan

⁴ Department of Earth and Space Science, Graduate School of Science, Osaka university, Osaka, 560-0043, Japan

(Dated: October 10, 2012)

The accurate calculation of long-term phase evolution of gravitational wave (GW) forms from extreme (intermediate) mass ratio inspirals (E(I)MRIs) is an inevitable step to extract information from this system. Achieving this goal, it is believed that we need to understand the gravitational self-forces. However, it is not quantitatively demonstrated that the second order self-forces are necessary for this purpose. In this paper we revisit the problem to estimate the order of magnitude of the dephasing caused by the second order self-forces on a small body in a quasi-circular orbit around a Kerr black hole, based on the knowledge of the post-Newtonian (PN) approximation and invoking the energy balance argument. In particular, we focus on the averaged dissipative part of the self-force, since it gives the leading order contribution among the various components of them. To avoid the possibility that the energy flux of GWs becomes negative, we propose a new simple resummation called *exponential resummation*, which assures the positivity of the energy flux. In order to estimate the magnitude of the yet unknown second order self-forces, here we point out the scaling property in the absolute value of the PN coefficients of the energy flux. Using these new tools, we evaluate the expected magnitude of dephasing. Our analysis indicates that the dephasing due to the second order self-forces for quasi-circular E(I)MRIs may be well captured by the 3PN energy flux, once we obtain all the spin dependent terms, except for the case with an extremely large spin of the central Kerr black.

PACS numbers: 04.30Db, 04.25Nx, 04.30Tv, 95.85.Sz

I. INTRODUCTION

Extreme mass ratio inspirals (EMRIs) and intermediate mass ratio inspirals (IMRIs), in which a stellar mass or several tens solar mass compact object inspirals into a more massive central black hole, have attracted much interest not only as a promising source of the gravitational waves (GWs) for future space born GW detectors, but also as a unique clean probe of the spacetime region of strong gravity. To achieve the test of general relativity using GWs from E(I)MRIs, we need to predict sufficiently accurate waveforms. This requirement motivates us to model the E(I)MRIs as the motion of a small body in a given background spacetime with gravitational back reaction. This back reaction is treated as gravitational self-forces [1–6] and its higher order extension with respect to the mass ratio has attracted much interest in recent years [7–11]. See following review articles, *e.g.*, [12, 13] and references therein for more details.

In fact, following the scaling argument (See *e.g.* [14, 15]), the phase of GWs from a particle whose orbital frequency sweeps a few orders of magnitude before the plunge can be expanded as

$$\Phi = \frac{M_{\text{BH}}}{\mu} \left[\Phi^{(0)} + \frac{\mu}{M_{\text{BH}}} \Phi^{(1)} + O\left(\frac{\mu^2}{M_{\text{BH}}^2}\right) \right], \quad (1)$$

where $\Phi^{(0)}$ and $\Phi^{(1)}$ are $O(1)$ quantities independent of μ , which is the mass of a small particle, and M_{BH} and a are the mass and the spin parameter of the Kerr black hole, respectively. ¹ On the one hand, $\Phi^{(0)}$ in Eq. (1), we only need the self-forces up to the first order time-averaged dissipative part. In addition, it has been long known that $\Phi^{(0)}$ can be computed with well established balance argument, which relates the first order time-averaged dissipative part of self-forces to the energy and angular momentum fluxes associated to the global killing vectors on background Kerr black hole [19]. Though the Carter constant is

*Electronic address: isoyama@yukawa.kyoto-u.ac.jp

†Electronic address: ruyichi.fujita@uib.es

‡Electronic address: sago@artsci.kyushu-u.ac.jp

§Electronic address: tagoshi@vega.ess.sci.osaka-u.ac.jp

¶Electronic address: tanaka@yukawa.kyoto-u.ac.jp

¹ Throughout this paper, we assume that the resonance is absent during its orbital evolution. See Refs. [15–18] for further discussion about resonant orbits.

not associated to any Killing vector and simple balance argument is not applicable, the way to computing long time averaged evolution of Carter constant has been already well established [20–23]. On the other hands, $\Phi^{(1)}$, which is referred to as the dephasing relative to $\Phi^{(0)}$, depends on two different components of the self-forces: the first order conservative part, and the averaged second order dissipative part.² The first sub-leading term $\Phi^{(1)}$ still can be important since potentially it may give the correction significantly greater than unity to the phase [24–26].

To obtain accurate waveform, there have been many works on self-forces. As for the first order conservative part, thanks to the recent massive development, we are now in part ready for practical computation with numerical implementation [27–31]. Particularly, in the case of Schwarzschild background the correction to the orbital frequencies [32] and $\Phi^{(1)}$ [33] have been already studied extensively. Even in the case of Kerr background the preliminary results of the self-forces in quasi-circular orbits have been reported [34].

By contrast, the averaged second order dissipative part of self-forces has been so far studied only at the formal level in the context of black hole perturbation [7–11]. It will require much more effort to establish the method to compute the second order dissipative part, especially in the case of Kerr background. Under such circumstances, a typical strategy to evaluate $\Phi^{(1)}$ is to make use of the standard post-Newtonian (PN) approximation, in which we assume slow motion of a satellite and its weak gravitational field. Based on the PN approximation, Huerta and Gair [35] evaluated the size of dephasing caused by the first order conservative self-forces and the averaged second order self-forces, picking up representative EMRIs in quasi-circular orbits on a Kerr black hole. The same dephasing was also discussed by Yunes *et al.* [36] using the effective one body formalism again for representative E(I)MRIs.

A naive expectation is that the PN approximation is not suitable for modelling the waveforms of E(I)MRIs, especially for a Kerr black hole with large spin. Typical E(I)MRIs in circular orbits spend the last few years of inspiral in the vicinity of the inner most stable circular orbit (ISCO). Since the ISCO radius reaches the event horizon as the spin of the Kerr black hole is increased to the extremal limit, the motion of the body becomes highly relativistic, exceeding the validity range of the standard PN approximation [37]. However, the things are not so trivial. The time spent near the ISCO becomes longer as we increase the mass ratio M_{BH}/μ , but in that case the mass of the large central black hole also becomes larger. Then, the total cycles of GWs become smaller for a given observation time. As a result, the correction due to higher order self-forces might be suppressed below the observational threshold, despite the loss of the accuracy of the PN approximation. Therefore, it is not so obvious if there are E(I)MRIs such that really require the notion of the second order self-forces.

The previous analyses mentioned above [35, 36] are focusing on the corrections coming from the self-forces at the currently available PN order and are limited to representative E(I)MRIs. To get an insight into whether the second order self-forces based on the black hole perturbation are really necessary to calculate the waveforms of quasi-circular E(I)MRIs or not, therefore, it would be useful to give an estimate of dephasing coming from the averaged second order dissipative self-forces, focusing on the yet unknown higher PN terms and surveying the whole parameter region of E(I)MRIs.

What we discuss in this paper is the adiabatic evolution of E(I)MRIs in quasi-circular orbits on a Kerr spacetime. Here the adiabatic evolution means an approximation in which the evolution of the orbital frequency is determined by the energy balance argument, *i.e.*, the rate of change of the total energy of the binary is equated to the energy flux emitted to the infinity.

To evaluate the order of magnitude of the yet unknown higher PN corrections, we need to rely on some extrapolation. For this purpose, we first introduce a simple new resummation of the energy flux, which we call the “exponential resummation”. When the spin of a Kerr black hole is large enough, the PN energy flux in the Taylor form can be negative outside the ISCO radius for some PN orders [38]. If this happens, the estimated total phase before the plunge diverges and the extrapolation to the higher PN order will not make sense. Our simple “exponential resummation” is the one that ensures the positivity of the energy flux.

As a PN input for the corrections at the next leading order in the mass ratio, the best one available so far is the 3.5PN energy flux of GWs [39, 40] with linear spin dependent terms up to 3PN order, which has been recently derived by Blanchet *et al.* [41]. To estimate the possible magnitude of the yet unknown higher order PN terms, we focus on a scaling property among the PN coefficients in the energy flux. Using the 8PN energy flux in the test particle limit [42, 43], we will show that the absolute values of the coefficients scale roughly as required from the convergence of the PN series up to the light ring radius. (This point was also discussed by Nakano *et al.* independently [44].) Since this scaling behavior is related to the PN convergence, despite the lack of the higher PN terms in the energy flux, we conjecture that the same scaling property will hold for the terms higher order in the mass ratio. Under this assumption, we estimate the order of magnitude of the unknown portion of the energy flux coming from higher PN terms at the next leading order in the mass ratio via “extrapolation”. Gathering the tools stated above, we will investigate the impact on the dephasing from the averaged dissipative second-order self-forces for various E(I)MRIs systematically. This is the main goal of this paper.

² Here the “dissipative” part means the self-forces that cause the time variation of the constants of motion, such as the energy, the angular momentum around the axis of symmetry and the Carter constant. The “conservative” part means the one giving the correction to the relation between the orbital frequencies and the constants of motion [15, 20]. The meaning of “time-averaged” is averaging over a sufficient long period compared to the time scale for the evolution of the phase difference between the oscillations in the radial and the zenith angle directions.

The remainder of this manuscript is organized as follows. In Sec. II we briefly review how the accumulated phase of GWs from adiabatic inspiral is calculated, based on the balance argument. In Sec. III, to cure the negative energy flux of GWs that appears in the truncated PN Taylor series expansion, we propose the exponential resummation that ensures the positivity of the energy flux. Sec. IV is dedicated to the study of the scaling property in the coefficients of the energy flux in the test particle limit that becomes manifest owing to the brand new 8PN energy flux [43]. In Sec. V, using the exponential resummation and the scaling property, we will estimate the dephasing coming from the yet unknown part of the second order self-forces for various E(I)MRIs. We find that the unknown non-linear spin dependent terms at the lower PN order in the energy flux dominate the unknown dephasing. We summarize our results and conclude in Sec. VI.

In this manuscript, we use geometrical units $G = c = 1$ and the sign convention of the metric is $(-, +, +, +)$. The coordinates (t, r, θ, ϕ) denote the Boyer-Lindquist coordinates of the Kerr black hole [45]. We frequently use the dimensionless spin defined by $q := a/M_{\text{BH}}$, and the symmetric mass ratio defined by $\nu := \mu M_{\text{BH}} / (M_{\text{BH}} + \mu)^2$.

II. THE ACCUMULATED PHASE OF THE GRAVITATIONAL WAVE FROM AN INSPIRALING BINARY

We consider a binary composed of a small satellite body with the rest mass μ in a quasi-circular orbit around a Kerr black hole with the mass M_{BH} and the spin parameter a . We neglect the effect of the spin of the satellite, which is negligibly small for E(I)MRIs though plays no negligible role in some situations [46, 47]. For a binary in a quasi-circular orbit, the accumulated phase of GWs is calculated as

$$\Phi := -2 \int_{x_{\text{ISCO}}}^{x_0} dx \frac{x^{3/2}}{M} \frac{E'(x)}{\dot{E}(x)}, \quad (2)$$

where E is the binding energy of the binary, $\dot{E} := dE/dt$ is the energy loss rate, and the prime denotes the differentiation with respect to x , which is a dimensionless orbital frequency defined by $x := (M\Omega)^{2/3}$ with the orbital frequency Ω and the total mass $M := M_{\text{BH}} + \mu$. Here E is supposed to depend only on x , neglecting the effect of variation of mass due to the energy absorbed by the black hole. Throughout the paper we totally neglect the effect of this time dependent mass variation. We choose the lower bound of the integral in Eq. (2), x_{ISCO} to the value of x at the inner most stable circular orbit (ISCO) and the upper bound x_0 to the one determined by the condition coming from the finite observation time. Throughout this paper, we adopt $t_{\text{obs}} = 1$ year as the observation time.

Our primary interest in this paper is the order of magnitude of the dephasing coming from the corrections at the next leading order in the mass ratio. To evaluate it, we calculate the difference between Φ with and without the higher order corrections. Checking the detectability of this difference by observations, the initial and the final frequencies should be kept unchanged when we evaluate it. Here, we fix both x_0 and x_{ISCO} to the values determined by the test particle limit. Namely, [48]

$$\begin{aligned} x_{\text{ISCO}} &:= \left(R_{\text{ISCO}}^{3/2} + q \right)^{-1/6} & R_{\text{ISCO}} &:= 3 + Z_2 \mp \sqrt{(3 - Z_1)(3 + Z_1 + 2Z_2)}, \\ Z_1 &:= 1 + (1 - q^2)^{1/3} [(1 + q)^{1/3} + (1 - q)^{1/3}], & Z_2 &:= \sqrt{3q^2 + Z_1^2}, \end{aligned} \quad (3)$$

where the upper (lower) sign in the second equation is chosen for the co (counter)-rotating case with $q > (<) 0$ and $R := r/M_{\text{BH}}$ represents the dimensionless radius. x_0 is determined by

$$1 \text{ (year)} = - \int_{x_{\text{ISCO}}}^{x_0} dx \frac{E'^{[0]}(x)}{\dot{E}^{[0]}(x)}, \quad (4)$$

where the superscript $^{[0]}$ means the leading order contribution in the mass ratio, i.e., the test particle limit, and

$$E^{[0]} = \mu \frac{R^{3/2} - 2R^{1/2} + q}{R^{3/4}(R^{3/2} - 3R^{1/2} + 2q)^{1/2}}, \quad \Omega := \frac{1}{M(R^{3/2} + q)}, \quad R = \left(\frac{1 - qx^{3/2}}{x^{3/2}} \right)^{2/3}. \quad (5)$$

As for the energy, we here adopted the binding energy of the binary instead of the energy directly related to the four momentum of the particle. It would be more natural to consider the latter in the context of the self-force calculation based on the black hole perturbation, but it is not gauge invariant. Hence, the former is more suitable for comparison with the PN calculation. These two energies are different at the next leading order in the mass ratio due to the presence of the gravitational field energy, but there must be one-to-one correspondence between them, once the gauge is completely fixed.

The energy balance argument tells us that the averaged loss of the total energy should be equal to the averaged total energy flux to the future null infinity because of the energy conservation (see e.g. Ref. [19, 49]). Hence, as long as we define the binding energy appropriately,

$$-\frac{dE}{dt} = \mathcal{L}, \quad (6)$$

holds, in the sense averaged over a sufficiently long time and neglecting the horizon absorption flux, where $\mathcal{L}$ is the energy flux emitted to the null infinity. Hence, we evaluate the phase by

$$\Phi[\mathcal{L}] := 2 \int_{x_{\text{ISCO}}}^{x_0} dx \frac{x^{3/2}}{M} \frac{E'^{[0]}(x)}{\mathcal{L}(x)}. \quad (7)$$

In the above equation, as we are interested in the dissipative corrections, we fixed E' to the expression in the test particle limit. We expand $\mathcal{L}$ as $\mathcal{L} = \mathcal{L}^{[0]} + \nu \mathcal{L}^{[1]} + O(\nu^2)$. To compute $\mathcal{L}^{[0]}$, we can use the Teukolsky formalism [50] and invoke the numerical code developed by Fujita and Tagoshi [51, 52].

The finite mass corrections at the next leading order, $\mathcal{L}^{[1]}$, are in part provided by the standard PN calculations. In the PN formalism, the energy flux of GWs emitted to the infinity from a quasi-circular binary is obtained up to 3.5PN order for the spin independent terms [39, 40] and up to 3PN order for the terms linear in spin when the spin vectors are parallel to the orbital axis [41]. In the present notation, truncated at $O(\nu)$, it is given by

$$\begin{aligned} \mathcal{L}_{n\text{PN}}^{\text{T}}(x, q) = & \frac{32}{5} x^5 \nu^2 \left[1 + x \left(-\frac{1247}{336} - \frac{35}{12} \nu \right) + x^{3/2} \left(4\pi - \frac{11}{4} q + \frac{17}{4} q \nu \right) + x^2 \left(-\frac{44711}{9072} + \frac{9271}{504} \nu \right) \right. \\ & + x^{5/2} \left(-\frac{8191}{672} \pi - \frac{59}{16} q + \nu \left\{ -\frac{583}{24} \pi + \frac{3749}{144} q \right\} \right) \\ & + x^3 \left(\frac{6643739519}{69854400} + \frac{16}{3} \pi^2 - \frac{1712}{105} \gamma_E - \frac{856}{105} \log(16x) - \frac{65}{6} \pi q + \left\{ -\frac{134543}{7776} + \frac{41}{48} \pi^2 + \frac{33}{2} \pi q \right\} \nu \right) \\ & \left. + x^{7/2} \left(-\frac{16285}{504} + \frac{214745}{1728} \nu \right) \pi + O(x^4, \nu^2, q^2, qx^{7/2}) \right], \quad (8) \end{aligned}$$

where $\gamma_E = 0.57721 \dots$ denotes Euler's constant. We will refer to the above expression for the energy flux truncated at the n th PN order³ as “the n PN Taylor flux”. We denote it by $\mathcal{L}_{n\text{PN}}^{\text{T}}$, and expand it like $\mathcal{L}_{n\text{PN}}^{\text{T}} := \mathcal{L}_{n\text{PN}}^{\text{T}[0]} + \nu \mathcal{L}_{n\text{PN}}^{\text{T}[1]} + \dots$ as before. We also frequently use “the normalized n PN flux” defined by

$$L_{n\text{PN}}^{\text{T}}(x, q) := \left(\frac{32}{5} \nu^2 x^5 \right)^{-1} \mathcal{L}_{n\text{PN}}^{\text{T}}(x, q). \quad (9)$$

From now on, we denote the PN flux solely composed of the known part as “the known n PN Taylor flux” and distinguishing it from the full PN flux by associating “ $\sim$ ” like $\tilde{\mathcal{L}}_{n\text{PN}}^{\text{T}}$. We also introduce the notation to denote the residual terms higher order in PN expansion and in spin by

$$\mathcal{L}_{>n\text{PN}}^{\text{T}[i]} := \mathcal{L}_{\text{full}}^{[i]} - \mathcal{L}_{n\text{PN}}^{\text{T}[i]}. \quad (10)$$

Then, $\tilde{\mathcal{L}}_{>3.5\text{PN}}^{\text{T}[1]}$ represents the yet unknown energy flux to be determined from the computation of the second order dissipative self-forces. Further we define $\tilde{\mathcal{L}}_{n\text{PN}}^{\text{T}[0]}$ and $\tilde{\mathcal{L}}_{>n\text{PN}}^{\text{T}[0]}$ by the terms in $\mathcal{L}^{[0]}$ that correspond to $\tilde{\mathcal{L}}_{n\text{PN}}^{\text{T}[1]}$ and $\tilde{\mathcal{L}}_{>n\text{PN}}^{\text{T}[1]}$, respectively. To be precise, we define $\tilde{\mathcal{L}}_{n\text{PN}}^{\text{T}[0]}$ by the sum of the spin independent terms up to 3.5PN order and the terms linear in the black hole spin up to 3PN order. $\tilde{\mathcal{L}}_{>n\text{PN}}^{\text{T}[0]}$ is the remaining terms at the leading order in the mass ratio, which also includes the non-linear spin dependent terms in all PN orders.⁴

III. A MANIFESTLY POSITIVE DEFINITE ENERGY FLUX: EXPONENTIAL RESUMMATION

In this section we propose a new simple resummation scheme, which can be easily applied if we just know the n PN Taylor flux.

A. Limitation of the Taylor flux revisited

It is easy to imagine that the PN Taylor flux is not so accurate when the orbital radius becomes small. Indeed, Zhang *et al.* [37] pointed out that the n PN Taylor flux in the test particle limit rapidly loses the accuracy around the ISCO radius. However, it is a

³ We refer to a term of $O(x^n)$ relative to the leading order as those of n PN order.

⁴ Here we do not distinguish the logarithmic term in 3PN Taylor flux in the test particle limit from the other 3PN terms. We confirmed that the results in the present paper do not change much even if we exclude this term from $\tilde{\mathcal{L}}_{n\text{PN}}^{\text{T}[0]}$.




FIG. 1: The normalized n PN Taylor flux in the test particle limit, $L_{n\text{PN}}^{\text{T}[0]}(x, q)$, for $q = 0.9$ up to $x_{\text{ISCO}}(0.9) := 0.78014\dots$. The horizontal axis is the dimensionless frequency x of a body. The label “Exact” means the exact numerical energy flux, calculated by Fujita and Tagoshi [51].

different issue whether or not the Taylor flux is accurate enough for our present purpose because what we are interested in here is whether or not the dephasing in GW waveforms is measurebale.

As we want to evaluate the correction to Φ in Eq. (7) caused by the yet unknown part of the flux $\tilde{\mathcal{L}}_{>n\text{PN}}^{\text{T}[1]}$, we need some extrapolation method. For this purpose, later it becomes necessary to evaluate something like the phase for the flux truncated at n PN order, $\Phi[\tilde{\mathcal{L}}_{n\text{PN}}^{\text{T}[0]}]$. At this point we find that the n PN Taylor flux is problematic. As mentioned earlier, the energy flux truncated at some PN orders become negative outside R_{ISCO} if the spin of black hole q is sufficiently large, firstly pointed out by Tagoshi *et al.* [38].

To see how serious the problem is, we revisit the Taylor flux in the test particle limit. The spin independent terms in $\mathcal{L}_{n\text{PN}}^{\text{T}[0]}$ have been analytically calculated up to 22PN order and the spin dependent terms up to 8PN order by Fujita [42, 43]. The expression of the normalized 8PN energy flux including complete spin dependent terms is schematically expanded as⁵

$$L_{8\text{PN}}^{\text{T}[0]}(x, q) = \sum_{n=0}^8 \sum_{p=0}^{p_{\text{max}}} \hat{L}^{(n,p)}(q) x^n (\log(x))^p, \quad (11)$$

where p_{max} is the maximum integer that does not exceed $n/3$. In FIG. 1, we depicted $L_{n\text{PN}}^{\text{T}[0]}(x, q)$, the normalized n PN Taylor flux in the test particle limit (11) truncated at various PN orders for $q = 0.9$. The curves terminate at the ISCO frequency. We also plot the exact numerical energy flux in the test particle limit as a reference. The exact numerical energy flux is manifestly positive definite for $x < x_{\text{ISCO}}(0.9)$, while all the n PN Taylor fluxes in FIG. 1 cross zero at some $x = x_0(q) < x_{\text{ISCO}}(0.9)$. In addition to the already known 2.5PN and 4PN cases⁶ [38], we also find that 5PN, 5.5PN, 6.5PN and 8PN fluxes become negative before x reaches $x_{\text{ISCO}}(q)$. Even for a moderate value such as $q = 0.7$, we still observe the flux to cross zero, say at 8PN order. Once this happens, the integrand of (7) diverges, and then the phase evaluated by using the truncated flux $\mathcal{L}_{n\text{PN}}^{\text{T}[0]}$ does not make sense. For a reliable extrapolation, it is necessary to use a resummed expression for the energy flux that gives at least a finite estimate of the phase for the truncation at any PN order.

⁵ The explicit expression for $\mathcal{L}_{8\text{PN}}^{\text{T}[0]}$ will be made available elsewhere in an appropriate form [43].

⁶ While Tagoshi *et al.* [38] reported that 3PN Taylor flux with $q = 0.9$ became negative for small radius, we find that the 3PN Taylor flux is always positive definite for all radius irrespective of the spin parameter of the Kerr black hole.

B. The exponential resummation and an improved hybrid energy flux

To overcome the drawback of the n PN Taylor flux, a naive requirement will be to guarantee the energy flux to be always positive by resummation. Among various known resummation techniques [53–56], to the best of our knowledge, only the factorized resummation ensures the positivity of the energy flux. This resummation was proposed by Damour *et al.* [57] for the equal mass non-spinning binaries in a circular orbit. Fujita and Iyer [58] and Pan *et al.* [59] applied it to the test particle in a circular orbit around a Schwarzschild and a Kerr black hole, respectively. Here, we propose another even simpler resummation, which is defined by

$$\mathcal{L}_{n\text{PN}}^{\text{exp}}(x, q) := \frac{32}{5} \nu^2 x^5 \exp[L_{n\text{PN}}^{\text{exp}}(x, q)], \quad (12)$$

with

$$L_{n\text{PN}}^{\text{exp}}(x, q) := \log \left[L_{n'\text{PN}}^{\text{T}}(x, q) \right] \Big|_{\text{truncated at } n\text{PN order}}, \quad (13)$$

where $n' \geq n$ is understood. We refer to this resummation and the energy flux (12) as the exponential resummation and “the n PN exponential resummed flux”, respectively. This flux is manifestly positive and can incorporate any higher order corrections if the counterparts in the n PN Taylor flux is given. Furthermore, it significantly improves the accuracy of approximation [43]. For definiteness, we explicitly write the definition of $\mathcal{L}_{n\text{PN}}^{\text{exp}[0]}$ and $\mathcal{L}_{n\text{PN}}^{\text{exp}[1]}$ as

$$\mathcal{L}_{n\text{PN}}^{\text{exp}[0]}(x, q) := \frac{32}{5} \nu^2 x^5 \exp[L_{n\text{PN}}^{\text{exp}[0]}(x, q)], \quad \mathcal{L}_{n\text{PN}}^{\text{exp}[1]}(x, q) := \left(\frac{32}{5} \nu^2 x^5 \exp[L_{n\text{PN}}^{\text{exp}[0]}(x, q)] \right) L_{n\text{PN}}^{\text{exp}[1]}(x, q), \quad (14)$$

with $L_{n\text{PN}}^{\text{exp}} = L_{n\text{PN}}^{\text{exp}[0]} + \nu L_{n\text{PN}}^{\text{exp}[1]} + \dots$. We show the explicit form only for $L_{n\text{PN}}^{\text{exp}[1]}$ as

$$\begin{aligned} L_{n\text{PN}}^{\text{exp}[1]}(x, q) = & -\frac{35}{12}x + \frac{17}{4}qx^{3/2} + \frac{30523}{4032}x^2 + \left(\frac{136229}{4032}q - \frac{101}{8}\pi \right) x^{5/2} \\ & + \left(-\frac{43670915}{12192768} + \frac{41}{48}\pi^2 - \frac{\pi}{2}q \right) x^3 + \frac{70075}{6048}\pi x^{7/2} + O(x^4, qx^{7/2}, q^2). \end{aligned} \quad (15)$$

Combining the exact numerical flux obtained in the test particle limit, $\mathcal{L}_{\text{full}}^{[0]}$, and the PN flux, $\mathcal{L}_{\text{PN}}$, one can obtain a better estimate for the energy flux. In the case of the Taylor flux, these two are combined simply by summing $\mathcal{L}_{\text{full}}^{[0]}$ and $\mathcal{L}_{\text{PN}}^{\text{T}[1]}$. In the case of the exponential resummed energy flux (12), we need to do this summation at the level of the exponent, L^{exp} . Namely, expanding $L_{n\text{PN}}^{\text{exp}}$ in powers of ν , we replace $L_{n\text{PN}}^{\text{exp}[0]}$ with the one corresponding to the numerical flux. Then, we obtain

$$\mathcal{L}_{n\text{PN}}^{\text{hyb}}(x, q) = \mathcal{L}_{\text{full}}^{[0]}(x, q) \exp \left[\nu L_{n\text{PN}}^{\text{exp}[1]}(x, q) + \dots \right], \quad (16)$$

which we call “the n PN hybrid flux”.

In the analogous way as before, we introduce “the known n PN exponential resummed flux” and “the known n PN hybrid flux”, and distinguish them from their respective counterparts by associating “ $\sim$ ”. Here the truncation for the known part is made at the level of $L^{\text{exp}[1]}$. Namely, $\tilde{L}_{n\text{PN}}^{\text{exp}[1]}$ is truncated at the same PN order for each order of spin that is included in the known terms $\tilde{L}_{n\text{PN}}^{\text{T}[1]}$. For the exponential resummed flux, we can also introduce $\tilde{L}_{n\text{PN}}^{\text{exp}[0]}$, the counterpart of $\tilde{L}_{n\text{PN}}^{\text{exp}[1]}$ in the test particle limit, as in the case of $\tilde{L}_{n\text{PN}}^{\text{T}[0]}$. Namely, $\tilde{L}_{n\text{PN}}^{\text{exp}[0]}$ is truncated at the same PN order for each order of spin that is included in the known terms $\tilde{L}_{n\text{PN}}^{\text{T}[1]}$. On the other hand, $\tilde{L}_{n\text{PN}}^{\text{hyb}[0]}$ does not make sense. In the following we identify the hybrid flux with the exponential resummed flux for the leading order in the mass ratio. We summarize our notation for various fluxes used in the rest of our paper in Table I, for readability.

IV. THE SCALING LAW OF THE COEFFICIENTS IN THE ENERGY FLUX

Now we tackle our main issue: how to evaluate the magnitude of the yet unknown part of the energy flux, $\tilde{\mathcal{L}}_{>3.5\text{PN}}^{[1]}$. (When we discuss general issues independent of the form of the energy flux, we simply suppress the labels “T”, “exp” and “hyb”.) Our strategy here is to establish the scaling property in the coefficients of the PN expansion of the energy flux in the test particle limit for fixed q . (See also Ref. [44].) The argument of the n PN exponential resummed flux, $L_{n\text{PN}}^{\text{exp}[0]}$, is expanded as

$$L_{n\text{PN}}^{\text{exp}[0]}(x, q) := \sum_n \left(\frac{x}{x_{\text{lr}}(q)} \right)^n \sum_p C_{n,p}(q) \left(\log \left[\frac{x}{x_{\text{lr}}(q)} \right] \right)^p, \quad (17)$$

	Symbol
The flux truncated at n PN order	$\mathcal{L}_{n\text{PN}}^{[i]}$
The residual part of $\mathcal{L}_{n\text{PN}}^{[i]}$	$\mathcal{L}_{>n\text{PN}}^{[i]}$
The known part of flux truncated at n PN order	$\tilde{\mathcal{L}}_{n\text{PN}}^{[i]}$
The residual of $\tilde{\mathcal{L}}_{n\text{PN}}^{[i]}$	$\tilde{\mathcal{L}}_{>n\text{PN}}^{[i]}$
The normalized n PN Taylor flux	$L_{n\text{PN}}^{\text{T}[i]}$
The exponent of n PN exponential resummed flux	$L_{n\text{PN}}^{\text{exp}[i]}$

TABLE I: Our notation for various fluxes in this paper. Upper index $[i]$ refers to the order of truncation with respect to the mass ratio ν : $\mathcal{L} = \mathcal{L}^{[0]} + \nu\mathcal{L}^{[1]} + O(\nu^2)$. We also put label “T”, “exp”, “hyb” and “full” which respectively correspond to Taylor, exponential, hybrid and numerical complete flux, to distinguish the flux type.




FIG. 2: The absolute value of the contribution to the energy flux from each PN order $|C_{n,0}|$ at $x = x_{\text{lr}}(q)$, with the horizontal axis being the PN order n . Note that the values of $|C_{n,0}|$ are accidentally small with some fixed q . We safely excluded these values from the plot.

where

$$x_{\text{lr}}(q) := \left(\frac{1}{R_{\text{lr}}^{3/2} + q} \right)^{2/3}, \quad (18)$$

with

$$R_{\text{lr}}(q) := 2 \left[1 + \cos \left(\frac{2}{3} \arccos q \right) \right], \quad (19)$$

is the value of x corresponding to the radius of the circular orbit on the light ring [48].

It is known that for circular orbits the source term of the Teukolsky equation has a simple pole when the orbit is on the light ring radius, *i.e.*, at $x = x_{\text{lr}}(q)$ [60, 61]. The light ring radius is the innermost radius for the presence of an (unstable) circular orbit, and there the particle energy per unit rest mass diverges. As a result, the energy flux also diverges there. Besides the light ring singularity, another origin of singularity in solving the Teukolsky equation is the presence of quasi-normal mode poles, which are zeros of the Wronskian. However, the absolute value of the first quasi-normal mode frequency, which is the smallest, is almost identical to the frequency of GWs from the source at the light ring [62]. Hence, we expect that x_{lr} determines, at least approximately, the convergence radius when the energy flux is seen as an analytic function of x . If the PN expansion of the energy flux converges as long as $x < x_{\text{lr}}(q)$, $x = x_{\text{lr}}(q)$ is the first place where the series ceases to converge.

With this expectation in mind, we plot the contribution from each PN order with $x = x_{\text{lr}}(q)$, $|C_{n,0}|$, in FIG. 2 for various values of q . Here $q = 0.998$, is a bound for possible maximum spin of an astrophysical black hole called Thorne limit [63]. As

is expected, $|C_{n,0}|$ is roughly independent of the PN order n , as typical at the boundary of convergence for $|q| < 0.9$. Contrary to our expectation, for $|q| > 0.9$, $|C_{n,0}|$ continues to increase up to 8PN, and the series seems to diverge. However, this does not mean that PN series does not converge for $x \approx x_{\text{lr}}(q)$ with $|q| \approx 1$. When we make a similar plot for each partial wave contribution for a higher multipole component, the contribution to the energy flux peaks around a higher PN order, but the series is convergent, although we cannot check extremely high multipole components. On the other hand, the summation over multipole components is known to converge from the numerical computation of the energy flux by Fujita and Tagoshi [51, 52]. These facts may indicate that the convergence depends on the order to take summation. The point of FIG. 2 is that $|C_{n,0}|$ shows a nice scaling property with respect to the PN order for any value of q . Even for q close to the extremal limit, the plot becomes flat if we substitute a little smaller value of x instead of x_{lr} .

Now we are in the position to discuss how to estimate the order of magnitude of the yet unknown part of the energy flux, $\tilde{\mathcal{L}}_{>3.5\text{PN}}^{[1]}$. The second order perturbed Einstein equation would schematically take the form $\square h^{(2)} = (T[z + \delta z] - T[z]) + h^{(1)}T[z] + (\nabla h^{(1)})^2$, where $h^{(1)}$ and $h^{(2)}$ are, respectively, the first and the second order perturbations induced by a body. Here z is the background Kerr geodesic and δz is the $O(\nu)$ correction to it [65]. We know that all the first order perturbations have singularity only at $x \approx x_{\text{lr}}(q)$. The Green's function $\square^{-1}$ is basically the same as in the linear case. Hence, even for the second order perturbation, the convergence radius in x will be the same as in the linear case, and thus a similar scaling for the PN coefficients will be expected for the next leading order in the mass ratio, too. Then, we can guess the amplitude of the higher PN coefficients from the first few terms in the PN expansion that are known from the standard PN calculation.

One may worry that $\square^{-1}$ has singularities at the quasi-normal mode frequencies. When we consider the second order perturbation, there may arise high frequency components which are absent at the level of linear perturbation. However, this might not happen. The point is that the metric perturbation caused by a circular geodesic has a helical Killing vector⁷. At the level of second order source term, this helical symmetry is broken due to the presence of the deviation from the geodesic, δz . Apart from this contribution, however, the source term keeps the helical symmetry and hence the frequency that appears in a partial wave labelled by (ℓ, m) is $m\Omega$ only. Namely, higher frequency modes do not arise. Even if we take into account δz , the time scale associated with δz is as slow as $\Omega^{-1} \times O(\nu^{-1})$. Therefore, the presence of δz alters the frequency $m\Omega$ only by a small amount of $O(\nu\Omega)$. This essentially does not change the convergence radius in x .

Even if we get the order of magnitude of the PN coefficients, the scaling property tells us nothing about the actual sign of each term. Therefore, when x is close to $x_{\text{lr}}(q)$, it becomes difficult to guess the order of the magnitude of the infinite summation. However, since the higher order PN terms are suppressed by the power of $x/x_{\text{lr}}(q)$, the summation is dominated by a few leading terms when $x/x_{\text{lr}}(q)$ is reasonably small. In that case, we do not have to worry about the infinite summation. Here, we bravely step a little forward by making the following proposal that the ratio between $\tilde{\mathcal{L}}_{n\text{PN}}^{[1]}$ and $\tilde{\mathcal{L}}_{>n\text{PN}}^{[1]}$ will be the same order as the ratio between $\tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}$ and $\tilde{\mathcal{L}}_{>n\text{PN}}^{[0]}$. Based on this assumption, we extrapolate the known results to estimate the unknown $\tilde{\mathcal{L}}_{>n\text{PN}}^{[1]}$.

V. THE DEPHASING DUE TO THE AVERAGED DISSIPATIVE PORTION OF THE SECOND ORDER SELF-FORCES

The main focus of this section is to evaluate the dephasing from the averaged dissipative part of the second order self-forces in E(I)MRIs, based on the idea of the extrapolation proposed in the preceding section. In this paper, we measure the magnitude of the dephasing due to higher order corrections to the energy flux $\delta\mathcal{L}$ by

$$\Delta\Phi[\mathcal{L}, \delta\mathcal{L}] := -2 \int_{x_{\text{ISCO}}}^{x_0} dx \frac{x^{3/2} E'^{[0]}(x)}{M} \left| \frac{1}{\mathcal{L}(x) + \delta\mathcal{L}(x)} - \frac{1}{\mathcal{L}(x)} \right|. \quad (20)$$

Here the absolute value of the difference of fluxes is taken in the integrand. The factor $(\mathcal{L} + \delta\mathcal{L})^{-1} - \mathcal{L}^{-1}$ can change its signature in the domain of the integral for some parameter region of E(I)MRIs. If we do not take the absolute value, therefore, there might be an accidental cancellation between positive and negative contributions. Then, the order of magnitude of the evaluated $\Delta\Phi$ can largely deviate from what we really want to measure. Notice that, even if $\Delta\Phi$ defined without taking the absolute value in the integrand strictly vanishes, the deviation in the GW waveform is still detectable. Therefore, to avoid this possible underestimate of the dephasing, we take the absolute value of the difference.

As a general remark, we would like to mention the mass dependences of $\Phi[\mathcal{L}_{\text{ref}}^{[0]}]$, the phase for a certain reference flux $\mathcal{L}_{\text{ref}}^{[0]}$ which includes the leading PN terms in the test particle limit, and $\Delta\Phi[\mathcal{L}_{\text{ref}}^{[0]}, \delta\mathcal{L}^{[0]}]$ and $\Delta\Phi[\mathcal{L}_{\text{ref}}^{[0]}, \delta\mathcal{L}^{[1]}]$, the dephasings due to

⁷ The helical Killing vector $t^\alpha + \Omega\phi^\alpha$, that remains to generate a symmetry even for the perturbed spacetime: where t^α and ϕ^α are the asymptotically time translation and rotational Killing vectors of the background Kerr black hole, respectively. See e.g. Ref. [64] for more details.




FIG. 3: The dephasing due to the higher PN corrections in the test particle limit, $\nu \Delta\Phi[\mathcal{L}_{n\text{PN}}^{\text{exp}[0]}, \mathcal{L}_{>n\text{PN}}^{\text{exp}[0]}]$ as a function of ν/M for $q = \pm 0.9$.

$\delta\mathcal{L}^{[0]}$ and $\delta\mathcal{L}^{[1]}$, which are some portions of the energy flux at the leading order and the next leading order in the mass ratio, respectively. From the integral in Eq. (4), which defines x_0 , one can factor out M_{BH}/ν since

$$E'^{[0]} \propto M_{\text{BH}}\nu, \quad \mathcal{L}_{\text{ref}}^{[0]} \propto \nu^2, \quad (21)$$

for a given x . Hence, we find that x_0 is a function of M_{BH}/ν . Substituting $\delta\mathcal{L}^{[0]} \propto \nu^2$ and $\delta\mathcal{L}^{[1]} \propto \nu^3$ together with the relations (21) into the definitions of $\nu \times \Phi[\mathcal{L}_{\text{ref}}^{[0]}]$, $\nu \times \Delta\Phi[\mathcal{L}_{\text{ref}}^{[0]}, \delta\mathcal{L}^{[0]}]$ and $\Delta\Phi[\mathcal{L}_{\text{ref}}^{[0]}, \delta\mathcal{L}^{[1]}]$, we find that their mass dependences remain only through x_0 . Therefore, one can conclude that $\nu \times \Phi[\mathcal{L}_{\text{ref}}^{[0]}]$, $\nu \times \Delta\Phi[\mathcal{L}_{\text{ref}}^{[0]}, \delta\mathcal{L}^{[0]}]$ and $\Delta\Phi[\mathcal{L}_{\text{ref}}^{[0]}, \delta\mathcal{L}^{[1]}]$ depend on M_{BH} and ν only through the combination M_{BH}/ν for a large mass ratio $\nu \ll 1$ for a fixed observation period before the plunge.

A. The error caused by the post-Newtonian truncation in the test particle limit

Before we evaluate the dephasing due to the yet unknown part of the energy flux at the next leading order in the mass ratio, we will assess the magnitude of the dephasing due to the PN truncation in the test particle limit $\Delta\Phi[\mathcal{L}_{n\text{PN}}^{[0]}, \mathcal{L}_{>n\text{PN}}^{[0]}]$ in this subsection and that coming from the known PN terms, $\Delta\Phi[\mathcal{L}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}]$, in the succeeding subsection.

Here, we study the quantity $\Delta\Phi[\mathcal{L}_{n\text{PN}}^{[0]}, \mathcal{L}_{>n\text{PN}}^{[0]}]$ to examine the PN convergence in terms of the phase error for various E(I)MRIs. We show the results only for the exponential resummed flux because the phase for the PN Taylor flux becomes ill-defined for several PN orders. (The hybrid flux is identical to the exponential resummed one by definition in the test particle limit.)

In FIG. 3 we plot $\nu \Delta\Phi[\mathcal{L}_{n\text{PN}}^{\text{exp}[0]}, \mathcal{L}_{>n\text{PN}}^{\text{exp}[0]}]$ as a function of ν/M for $q = \pm 0.9$. The main message of these plots is that the PN convergence is almost uniform except for the case of large black hole spin for large ν/M . When q is large, ISCO becomes close to the light ring radius where PN convergence becomes very slow. Even in that case, for lower PN orders the convergence is rather smooth for large ν/M . This is because up to 2PN order the phase correction is dominated by the lower bound of the x -integral in (20), if the initial separation of the binary at the time 1 year before the plunge is sufficiently large. The reason why we focus on large ν/M is, as will be discussed in Append. V, the dephasing from the known PN terms is already well suppressed below 1 rad if ν/M is small enough. Thus, the PN convergence for small ν/M does not affect the following discussion.

Notice also that x_0 becomes smaller and smaller for larger ν/M . The 2.5PN case is marginal in the sense that all range of x almost equally contributes. By contrast, the contribution near the ISCO dominates for the corrections at the 3PN order or higher. Although the PN convergence is not clearly seen for $q > 0.9$, we should note that the phase error is never extremely enhanced for some particular post-Newtonian order, largely exceeding the values for 3PN or 3.5PN order, which we mainly focus on in the following discussion.

B. The expected dephasing from the unknown higher order post-Newtonian terms

Now we move on to our main issue: evaluating the dephasing due to the yet unknown higher order PN terms, $\Delta\Phi[\mathcal{L}_{\text{full}}^{[0]} + \tilde{\mathcal{L}}_{3.5\text{PN}}^{[1]}, \tilde{\mathcal{L}}_{>3.5\text{PN}}^{[1]}]$, based on the idea for the extrapolation proposed in Sec. IV. The scaling argument will tell us that the dephas-




FIG. 4: The expected residual dephasing $\Delta\Phi_{\text{guess}}[\mathcal{L}_{\text{full}}^{[0]} + \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}, \tilde{\mathcal{L}}_{>n\text{PN}}^{[1]}]$ caused by the unknown part of the averaged dissipative second order self-forces. For $n = 3$, the 3.5PN known spin independent part of the flux at the next leading order in the mass ratio is treated as unknown.

ings coming from the respective PN terms at the leading order in the mass ratio will be roughly proportional to the corresponding next leading order dephasings. Namely,

$$\Delta\Phi[\mathcal{L}_{0\text{PN}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}] : \Delta\Phi[\tilde{\mathcal{L}}_{n\text{PN}}^{[0]}, \tilde{\mathcal{L}}_{>n\text{PN}}^{[0]}] \approx \Delta\Phi[\tilde{\mathcal{L}}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}] : \Delta\Phi[\mathcal{L}_{\text{full}}^{[0]} + \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}, \tilde{\mathcal{L}}_{>n\text{PN}}^{[1]}], \quad (22)$$

will hold. Recall that $\tilde{\mathcal{L}}_{n\text{PN}}^{[0]}$ consists of the terms at the leading order in mass ratio up to the same $n\text{PN}$ order that is available in $\tilde{\mathcal{L}}_{n\text{PN}}^{[1]}$.

Here, we have basically three options in the choice of the flux among the Taylor flux, the exponential resummed flux and the hybrid flux. (Recall that the hybrid flux is defined as such that is identical to the exponential resummed flux at the leading order in the mass ratio.) On one hand, the Taylor flux is problematic since it can be negative before reaching ISCO for some PN orders, say 2.5PN, with modelate value of the spin parameter as we mentioned earlier. In such cases the phase before the plunge is not well defined. On the other hand, the difference between the exponential resummed flux and the hybrid flux is negligible. Therefore, we here consider the hybrid flux only.

The residual dephasing estimated by using (22)

$$\Delta\Phi_{\text{guess}}[\mathcal{L}_{\text{full}}^{[0]} + \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}, \tilde{\mathcal{L}}_{>n\text{PN}}^{[1]}] := \frac{\Delta\Phi[\tilde{\mathcal{L}}_{n\text{PN}}^{[0]}, \tilde{\mathcal{L}}_{>n\text{PN}}^{[0]}]}{\Delta\Phi[\mathcal{L}_{0\text{PN}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}]} \Delta\Phi[\tilde{\mathcal{L}}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}], \quad (23)$$

with $n = 3$ and $n = 3.5$ is depicted in FIG. 4. The trend of the estimated dephasing is independent of the truncated PN order. The point of FIG. 4 is that the residual dephasing is rather suppressed over the whole range of the binary parameters. Indeed, the residual dephasing is at most about 10 rad in the case of 3PN truncation. In the case of 3.5PN truncation, the maximum value of the residual dephasing is bigger by factor 2 or so. Since the plot is given in the unit of radian, the value must be divided by 2π to translate it into the number of cycles. From the above results, one may say that the residual dephasing due to the yet unknown PN corrections at the next leading order in the mass ratio, $\Delta\Phi_{\text{guess}}[\mathcal{L}_{\text{full}}^{[0]} + \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}, \tilde{\mathcal{L}}_{>n\text{PN}}^{[1]}]$ is not negligible, adopting one cycle as the criteria for the significant dephasing.

In FIG. 4, we see a tiny bump around $\nu M_{\odot}/M \approx 10^{-10} - 10^{-11}$ for some values of q . The reason why this bump appears at this position can be understood from the observation that $\tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}$ crosses zero within the domain of the integral, $x_0 < x < x_{\text{ISCO}}$ when the bump appears. When $\tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}$ crosses zero, the factor in the denominator of Eq. (23) $\Delta\Phi[\mathcal{L}_{0\text{PN}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}]$ suffers from more or less accidental suppression even though we take the absolute value in the integrand of $\Delta\Phi$ in (20). We denote the value of x at which the flux $\tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}$ crosses zero by x_{cross} . Then, if $x_0 \ll x_{\text{cross}}$ or $x_0 \gg x_{\text{cross}}$, this suppression does not produce much effect on the estimate of $\Delta\Phi[\mathcal{L}_{0\text{PN}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}]$. Therefore, the suppression becomes significant only for $x_0 \approx x_{\text{cross}}$. The value of x_{cross} is rather close to the value at ISCO, x_{ISCO} , but typically not extremely close to it. Hence, $x_0 \approx x_{\text{cross}}$ occurs when x_0 is neither extremely close to ISCO nor very small like $x_0 \ll 1$, which corresponds to the parameter region $\nu M_{\odot}/M \approx 10^{-10} - 10^{-11}$. Since now we find that this tiny bump is to be attributed to an accidental zero in $\tilde{\mathcal{L}}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}$, this bump would be regarded as an artificial feature. If it is far to remove the bumps from FIG. 4, we will find that the significantly large dephasing (in the sense of exceeding 1 cycle) will be expected only for $\nu M_{\odot}/M \gtrsim 10^{-9}$.




FIG. 5: The expected residual dephasing $\Delta\Phi_{\text{guess}}[\mathcal{L}_{\text{full}}^{[0]} + \mathcal{L}_{n\text{PN}}^{[1]}, \mathcal{L}_{>n\text{PN}}^{[1]}]$ when we assume that the spin dependent terms at the lower PN orders are all known.

C. Unknown dephasing expected when we know the lower PN order non-linear spin dependent terms in the energy flux

In the preceding subsection, we find that the dephasing caused by the unknown PN higher order terms at the next leading order in mass ratio can be $O(10)$ rad or more. However, these unknown terms in the flux include non-linear spin terms at 3PN or lower order and all spin dependent terms at 3.5PN order, and hence we excluded the corresponding terms from $\tilde{\mathcal{L}}_{n\text{PN}}^{[0]}$, which appears in the expression (23). In FIG. 4, relatively large dephasing is observed for large $|q|$, for which the non-linear spin terms will be important.

Here we consider how the estimate presented in the preceding subsection is modified once we obtain all the spin dependent terms up to 3PN or 3.5PN order. What we need to evaluate is the expression obtained by removing “ $\sim$ ” from (23), i.e.,

$$\Delta\Phi_{\text{guess}}[\mathcal{L}_{\text{full}}^{[0]} + \mathcal{L}_{n\text{PN}}^{[1]}, \mathcal{L}_{>n\text{PN}}^{[1]}] := \frac{\Delta\Phi[\mathcal{L}_{n\text{PN}}^{[0]}, \mathcal{L}_{>n\text{PN}}^{[0]}]}{\Delta\Phi[\mathcal{L}_{0\text{PN}}^{[0]}, \mathcal{L}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}]} \Delta\Phi[\mathcal{L}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}]. \quad (24)$$

Here, one may think that $\tilde{\mathcal{L}}_{n\text{PN}}^{[1]}$ in the factor $\Delta\Phi[\mathcal{L}_{n\text{PN}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{[1]}]$ should have been also replaced with $\mathcal{L}_{n\text{PN}}^{[1]}$. Since we do not have the expression for $\mathcal{L}_{n\text{PN}}^{[1]}$ at hand, we cannot perform this replacement. However, the difference between $\mathcal{L}_{n\text{PN}}^{[1]}$ and $\tilde{\mathcal{L}}_{n\text{PN}}^{[1]}$ will not be significant because their dominant PN terms are common.

The resultant dephasing $\Delta\Phi_{\text{guess}}[\mathcal{L}_{\text{full}}^{[0]} + \mathcal{L}_{n\text{PN}}^{[1]}, \mathcal{L}_{>n\text{PN}}^{[1]}]$ is depicted in FIG. 5. In contrast to the FIG. 4, except for the near extremal spin case ($q = 0.998$) with the 3.5PN truncation ($n = 3.5$), the residual dephasing for large ν/M evaluated by using the hybrid flux does not exceed 1 rad. The residual dephasing exceeds 1 rad only at around the peak, which is likely to be an artifact due to the suppression in $\Delta\Phi[\mathcal{L}_{0\text{PN}}^{[0]}, \mathcal{L}_{n\text{PN}}^{[0]} - \mathcal{L}_{0\text{PN}}^{[0]}]$ similarly to the previous case. The above result indicates that the dominant source of the error caused by the unknown part of the flux is in the spin dependent terms at the lower PN orders up to 3PN or 3.5PN order. Namely, the knowledge of non-linear spin dependent terms in the energy flux is crucial for reducing the uncontrolled dephasing caused by the averaged dissipative second order self-forces. In addition, once we obtain these spin dependent terms of the energy flux, the exponential resummation will improve the accuracy of it to the level almost sufficient for detection of E(I)MRIs in the whole interesting parameter region.

VI. SUMMARY AND CONCLUSION

To evaluate the second order self-forces in the context of the black hole perturbation is a challenging issue motivated by the goal to extract information from extreme (intermediate) mass ratio inspirals (E(I)MRIs). In this paper we have assessed the dephasing of the GW waveform from quasi-circular E(I)MRIs, caused by the averaged dissipative part of the second order self-forces, focusing on giving an order of magnitude estimate of the influence of the yet unknown higher order post Newtonian (PN) corrections. Using the balance argument, the dephasing is related to the correction to the emitted energy flux. Although it will still require further efforts to calculate the second order self-forces using the black hole perturbation, they are partially already known from the standard PN calculation. We gave an estimate on how large dephasing will be caused by the yet unknown higher order PN terms for the last 1 year of inspiral before the plunge, exhaustively exploring the whole possible parameter region of E(I)MRIs.

To give a guess for the yet unknown higher order PN terms, we first introduced a simple resummation method for the energy flux, which we call the exponential resummation. This is simply obtained by exponentiating the energy flux and truncating it at the known PN order in the exponent. This resummation has three merits: it ensures the positivity of the energy flux, it accelerates the PN convergence, and it can be applicable if we just know the PN Taylor flux. Since the PN Taylor flux in the test particle limit with large q is known to be negative outside the innermost stable circular orbit (ISCO) at several PN orders, the total phase before the plunge becomes ill-defined. Hence, we cannot discuss the amplitude of the higher PN corrections by using the PN Taylor flux. The exponential resummation solves this problem and improves the PN convergence in terms of the total phase before the plunge. When we discuss the finite mass corrections to the energy flux, we can combine the idea of the exponential resummed flux with the known exact energy flux in the test particle limit. We also proposed such a phenomenological energy flux, which we call the hybrid flux.

To examine the dephasing from the unknown part of the averaged dissipative second order self-forces, we need some extrapolation. Using the brand new 8PN energy flux in the test particle limit [43] that includes all spin dependent terms, we discovered that the order of the magnitude of the absolute values of the coefficients up to 8PN order approximately follows a simple scaling law, which is what we expect when the actual convergence radius of the PN expansion is at the light ring radius, irrespective of the value of dimensionless spin parameter q . Since there is no reason to expect that the energy flux diverges outside the light ring radius even if we take into account the finite mass corrections for quasi-circular E(I)MRIs, we are motivated to assume that the ratio between the magnitude of the terms at $O(\nu^0)$ and $O(\nu)$ at the same PN order will roughly stay independent of the PN order.

Based on this assumption, we estimated the unknown portion of the energy flux that comes from the higher PN terms at the next leading order in the mass ratio and evaluated the residual dephasing due to them. We find that the residual dephasing may exceed one cycle for $\nu M_\odot/M \gtrsim 10^{-10}$ and the spin of the Kerr blackhole $|q| > 0.5$, assuming 1 year observation period. For some parameters, we found a little enhancement of the estimated dephasing at $\nu M_\odot/M \approx 10^{-10} - 10^{-11}$. Since this enhancement is likely to be attributed to a mild accidental cancellation in the factor in the denominator of the estimator of the residual dephasing that we adopted, it might be fair to remove the bump that arises for this reason. Then, the residual dephasing exceeds one cycle only for relatively large $O(\nu/M)$ with $|q| > 0.5$. Even for rather extreme case like $q = 0.998$ (Thorne limit), the expected residual dephasing is at most a few cycles or so.

In the unknown flux, non-linear spin terms at lower PN orders are also included since they are not yet calculated in the context of PN approximation even at the leading order in the PN expansion. As large residual dephasing is expected only for large $|q|$, one may suspect that the residual dephasing might be dramatically reduced once we know all the non-linear spin dependent terms up to 3PN or 3.5PN order. Therefore we made analogous plots for the expected residual dephasing assuming that the non-linear spin dependent terms are also available. We found that the expected residual dephasing is further suppressed to be less than 1 rad for $|q| < 0.9$, as we expected. Even in the extreme case with $q = 0.998$, the expected dephasing becomes less than 1 rad if we use 3PN truncation to evaluate the residual. We think the reason why 3.5PN truncation is worse is simply because, for large q close to unity, ISCO is very close to the light ring radius and hence the PN convergence is very poor. Thus, the resulting estimate of the residual also fluctuates by a large order of magnitude in this limit. However, even if we rely on the 3.5PN truncation, the expected dephasing is at most about 1 cycle for $q = 0.998$.

Although one year observation period is assumed in all the above estimates, the results do not depend much on it. The change in the observation time varies the relation between x_0 and ν/M , and hence the curves in FIGs. 4 and 5 are shifted just horizontally leftward. The expected dephasing becomes large only for large ν/M , but x_0 is already sufficiently small in this case. Therefore the effect of changing the observation period for large ν/M appears only in FIG. 4, in which lower order PN terms are contributing. In fact, one can see that the curves in FIGs. 5 are already saturated for large ν/M .

To conclude, the residual dephasing caused by the unknown averaged dissipative part of the second order self-forces is estimated to be at most a few cycles. This dephasing is mostly to be attributed to the lower PN order non-linear spin dependent terms. If the PN expansion to the 3PN or 3.5PN order is completed including the dependence on the black hole spin, the residual is expected to be further reduced.

As a final remark, we should note that there is a possibility that we might be underestimating the residual dephasing by an order of magnitude, since the PN convergence is not very smooth, especially for positive large q . Even though our estimate turns out to be a good approximation to the real magnitude, we still have a chance to have a golden event with a large signal to noise ratio. In that case, one cycle or even one radian may not be a sufficient accuracy that is required for the template waveforms. Then, the dissipative part of the second order self-forces that cannot be captured by the PN expansion becomes necessary to extract the best physics from the observational data. Also we should mention that we focused on quasi-circular E(I)MRIs in this paper, but our scaling argument will not apply any more for significantly eccentric orbits, since there are many modes whose frequency exceeds the naive PN convergence radius that will be given by the absolute value of the first complex quasi-normal mode frequency. Therefore, the hybrid energy flux cannot be expected to remain a good approximation near the plunge for such systems. Hence, our analysis does not at all discourage the study on the averaged dissipative part of the second order self-forces. Our claim is that it will be possible to perform nearly the best analyses of most of (nearly) quasi-circular E(I)MRIs, without waiting for the full development of our knowledge about the averaged dissipative part of the second order self-forces.




FIG. 6: The dephasing caused by the known terms up to the 3PN order at the next leading order in the mass ratio, $\Delta\Phi_{\text{PN}}[\mathcal{L}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{3\text{PN}}^{\text{hyb}[1]}]$, for various dimensionless spin parameters q of a Kerr black hole, based on the 3PN hybrid flux (16).

Acknowledgments

SI acknowledges the support from the Grant-in Aid for JSPS Fellows 24-4281. RF is grateful for the support of the European Union FEDER funds, the Spanish Ministry of Economy and Competitiveness (projects FPA2010-16495 and CSD2007-00042) and the Conselleria d'Economia Hisenda i Innovacio of the Govern de les Illes Balears. RF also appreciate the warm hospitality at Yukawa Institute for Theoretical Physics where part of this work was completed. HT is supported by the Grand-in-Aid for Scientific Research (No 23540309) and TT is supported by the Grand-in-Aid for Scientific Research (Nos. 21111006, 21244033, 24103001 and 24103006). Finally, this work was supported by the Grant-in-Aid for the Global COE Program "The Next Generation of Physics, Spun from Universality and Emergence" from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Appendix A: Dephasing from the known post-Newtonian terms

This Appendix is dedicated to see the dephasing from the available PN terms in the next leading order in the mass ratio. In the bulk of our paper, we introduced three different types of PN flux formulas, Taylor, exponential resummed and hybrid, but there is no qualitative difference among $\Delta\Phi_{\text{PN}}[\mathcal{L}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{\text{T}[1]}]$, $\Delta\Phi_{\text{PN}}[\mathcal{L}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{\text{exp}[1]}]$ and $\Delta\Phi_{\text{PN}}[\mathcal{L}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{n\text{PN}}^{\text{hyb}[1]}]$, as long as we choose the exact numerical flux $\mathcal{L}_{\text{full}}^{[0]}$ as the reference flux. This is because the significant difference among various fluxes appears only in the higher PN residual parts such that we discussed in the Sec. V. Therefore, we only show $\Delta\Phi_{\text{PN}}[\mathcal{L}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{3\text{PN}}^{\text{hyb}[1]}]$ in FIG. 6. We stress that what we show is the dephasing caused by the terms not at the 3PN order but up to the 3PN order. The main message of FIG. 6 is that the corrections to the phase of GWs due to the terms higher order in the mass ratio are suppressed below 1 rad, even if we include the leading order of the PN expansion, for a binary with $\nu M_{\odot}/M < 10^{-12}$, say, when the mass of the central Kerr black hole is greater than $3 \times 10^6 M_{\odot}$ with the satellite mass fixed to $10 M_{\odot}$. This holds irrespective of the value of the dimensionless spin parameter q . This plot clearly denies the naive statement: the effects of higher PN corrections should be important for a more massive central black hole because the satellite stays near the ISCO for a longer period.

As for the results in literature to compare, for the binary with $M_{\text{BH}} := 10^6 M_{\odot}$, $q = 0$ and $\nu = 10^{-5}$, Heurte and Gair [35] reported 1.5 rad dephasing after the last one year of inspiral based on the 2PN Taylor flux, while Yunes *et al.* [36] reported 0.71 rad dephasing after the last 11.5 months of inspiral for the same binary parameters. Following our definition of the dephasing, we calculate $\Delta\Phi_{\text{PN}}[\mathcal{L}_{\text{full}}^{[0]}, \tilde{\mathcal{L}}_{2\text{PN}}^{\text{hyb}[1]}]$ to find it 1.46 rad, which we think is in a good agreement with the previous results within the variance due to the different definition of dephasing.




FIG. 7: The dephasing $\nu\Delta\Phi[\mathcal{L}_{\text{full}}^{[0,H]} + \mathcal{L}_{2.5\text{PN}}^{[0,H]}, \mathcal{L}_{>2.5\text{PN}}^{[0,H]}]$ due to the horizon absorption flux and the dephasing $\nu\Delta\Phi[\mathcal{L}_{n\text{PN}}^{\text{exp}[0]}, \mathcal{L}_{>n\text{PN}}^{\text{exp}[0]}]$ due to the higher PN corrections, for $n = 3$ and $n = 3.5$, and $q = \pm 0.9$

Appendix B: Dephasing due to the horizon absorption flux

In the bulk of our manuscript, we neglected the flux absorbed by the horizon. In the test particle limit, Tagoshi *et al.* [66] and Yunes *et al.* [36] have already shown that the energy flux absorbed through the horizon cannot be neglected, leading to a large dephasing, especially when the Kerr black hole has large spin. However, this does not mean that the corrections due to the horizon absorption flux at the next leading order in the mass ratio are also important. Naively the standard PN formalism is not suitable for calculating the horizon absorption flux since the black hole horizon is beyond the reach of the standard PN expansion, although there is a direction to evaluate the horizon absorption flux by relating it with the tidal field around each black hole [67, 68].

In this appendix, we shortly address the dephasing caused by the flux absorbed through the horizon in the test particle limit to see what can be said on the dephasing due to the horizon absorption flux at the next leading order in the mass ratio. The flux absorbed through the horizon in the test particle limit is expanded as [66]

$$\mathcal{L}_{n\text{PN}}^{[0,H]}(x) := \frac{32}{5}\nu^2 x^5 \left\{ \left(-\frac{1}{4}q - \frac{3}{4}q^3 \right) x^{5/2} + \left(-q - \frac{33}{16}q^3 \right) x^{7/2} + O(x^4) \right\}, \quad (\text{B1})$$

and the exact numerical flux is also calculable [51]. We denote the contribution from the first term in the curly brackets in the above expression by $\mathcal{L}_{2.5\text{PN}}^{[0,H]}$ and the remaining part of the horizon flux by $\mathcal{L}_{>2.5\text{PN}}^{[0,H]}$, in a similar manner as before. As we mentioned above, the horizon absorption flux at the next leading order in the mass ratio is yet under development, but the part that requires the second order metric perturbation should start with 1PN order higher than the leading terms in the test particle limit, starting at 3.5PN order. Hence, we adopt $\mathcal{L}_{>2.5\text{PN}}^{[0,H]}$, excluding $\mathcal{L}_{2.5\text{PN}}^{[0,H]}$, as the counterpart of the unknown horizon absorption flux at the next leading order in the mass ratio. Under this consideration, we compare the dephasing due to the horizon absorption flux

$$\Delta\Phi[\mathcal{L}_{\text{full}}^{[0]} + \mathcal{L}_{2.5\text{PN}}^{[0,H]}, \mathcal{L}_{>2.5\text{PN}}^{[0,H]}] \quad (\text{B2})$$

to that coming from the PN truncation $\Delta\Phi[\mathcal{L}_{n\text{PN}}^{\text{exp}[0]}, \mathcal{L}_{>n\text{PN}}^{\text{exp}[0]}]$ for $n = 3$ or $n = 3.5$ in the test particle limit. The results are plotted in FIG. 7 for $q = \pm 0.9$.

FIG. 7 indicates that $\nu\Delta\Phi[\mathcal{L}_{\text{full}}^{[0]} + \mathcal{L}_{2.5\text{PN}}^{[0,H]}, \mathcal{L}_{>2.5\text{PN}}^{[0,H]}]$ stays at most $O(10^{-1})$ rad for all E(I)MRI parameter region. Indeed, it is, at most, about 1/10 of $\nu\Delta\Phi[\mathcal{L}_{n\text{PN}}^{\text{exp}[0]}, \mathcal{L}_{>n\text{PN}}^{\text{exp}[0]}]$. For large q , the smallness of the effect of the horizon flux relative to the higher order PN corrections can be understood from the work by Hughes [69], who numerically found that the energy flux absorbed by a Kerr black hole is at best 10 times smaller than the one emitted to the infinity, with the aid of the fact that the PN expansion shows a poor convergence near the ISCO for large q . Thus, we expect that the residual dephasing due to the horizon absorption

flux will be minor compared with the higher PN flux to the infinity, also in the next leading order in the mass ratio.

-
- [1] Y. Mino, M. Sasaki and T. Tanaka, Phys. Rev. D **55**, 3457 (1997) [gr-qc/9606018].
 - [2] T. C. Quinn and R. M. Wald, Phys. Rev. D **56**, 3381 (1997) [gr-qc/9610053].
 - [3] C. R. Galley, B. L. Hu and S. -Y. Lin, Phys. Rev. D **74**, 024017 (2006) [gr-qc/0603099].
 - [4] C. R. Galley and B. L. Hu, Phys. Rev. D **79**, 064002 (2009) [arXiv:0801.0900 [gr-qc]].
 - [5] S. E. Gralla and R. M. Wald, Class. Quant. Grav. **25**, 205009 (2008) [Erratum-ibid. **28**, 159501 (2011)] [arXiv:0806.3293 [gr-qc]].
 - [6] A. Pound, Phys. Rev. D **81**, 024023 (2010) [arXiv:0907.5197 [gr-qc]].
 - [7] E. Rosenthal, Phys. Rev. D **72**, 121503 (2005) [gr-qc/0508050]. E. Rosenthal, Phys. Rev. D **73**, 044034 (2006) [gr-qc/0602066]. E. Rosenthal, Phys. Rev. D **74**, 084018 (2006) [gr-qc/0609069].
 - [8] S. Detweiler, Phys. Rev. D **85**, 044048 (2012) [arXiv:1107.2098 [gr-qc]].
 - [9] A. Pound, Phys. Rev. Lett. **109**, 051101 (2012) [arXiv:1201.5089 [gr-qc]].
 - [10] S. E. Gralla, Phys. Rev. D **85**, 124011 (2012) [arXiv:1203.3189 [gr-qc]].
 - [11] A. Pound, arXiv:1206.6538 [gr-qc].
 - [12] L. Barack, Class. Quant. Grav. **26**, 213001 (2009) [arXiv:0908.1664 [gr-qc]].
 - [13] E. Poisson, A. Pound and I. Vega, Living Rev. Rel. **14**, 7 (2011) [arXiv:1102.0529 [gr-qc]].
 - [14] T. Hinderer and E. E. Flanagan, Phys. Rev. D **78**, 064028 (2008) [arXiv:0805.3337 [gr-qc]].
 - [15] T. Tanaka, Prog. Theor. Phys. Suppl. **163**, 120 (2006) [gr-qc/0508114].
 - [16] Y. Mino, Prog. Theor. Phys. **113**, 733 (2005) [gr-qc/0506003].
 - [17] E. E. Flanagan and T. Hinderer, arXiv:1009.4923 [gr-qc].
 - [18] E. E. Flanagan, S. A. Hughes and U. Ruangsri, arXiv:1208.3906 [gr-qc].
 - [19] D. V. Gal'tsov, J. Phys. A **15**, 3737 (1982).
 - [20] Y. Mino, Phys. Rev. D **67**, 084027 (2003) [gr-qc/0302075].
 - [21] N. Sago, T. Tanaka, W. Hikida and H. Nakano, Prog. Theor. Phys. **114**, 509 (2005) [gr-qc/0506092].
 - [22] N. Sago, T. Tanaka, W. Hikida, K. Ganz and H. Nakano, Prog. Theor. Phys. **115**, 873 (2006) [gr-qc/0511151].
 - [23] S. Drasco and S. A. Hughes, Phys. Rev. D **73**, 024027 (2006) [gr-qc/0509101].
 - [24] L. M. Burko, Phys. Rev. D **67**, 084001 (2003) [gr-qc/0208034].
L. M. Burko, Class. Quant. Grav. **22**, S847 (2005).
 - [25] J. R. Gair, Class. Quant. Grav. **26**, 094034 (2009) [arXiv:0811.0188 [gr-qc]].
 - [26] J. Thornburg, arXiv:1006.3788 [gr-qc].
J. Thornburg, arXiv:1102.2857 [gr-qc].
 - [27] L. Barack and A. Ori, Phys. Rev. D **61**, 061502 (2000) [gr-qc/9912010].
 - [28] L. Barack, Y. Mino, H. Nakano, A. Ori and M. Sasaki, Phys. Rev. Lett. **88**, 091101 (2002) [gr-qc/0111001].
 - [29] Y. Mino, H. Nakano and M. Sasaki, Prog. Theor. Phys. **108**, 1039 (2003) [gr-qc/0111074].
 - [30] S. L. Detweiler and B. F. Whiting, Phys. Rev. D **67**, 024025 (2003) [gr-qc/0202086].
 - [31] L. Barack and A. Ori, Phys. Rev. D **67**, 024029 (2003) [gr-qc/0209072].
 - [32] N. Warburton, S. Akcay, L. Barack, J. R. Gair and N. Sago, Phys. Rev. D **85**, 061501 (2012) [arXiv:1111.6908 [gr-qc]].
 - [33] K. A. Lackeos and L. M. Burko, arXiv:1206.1452 [gr-qc].
 - [34] A. G. Shah, J. L. Friedman and T. S. Keidl, arXiv:1207.5595 [gr-qc].
 - [35] E. A. Huerta and J. R. Gair, Phys. Rev. D **79**, 084021 (2009) [Erratum-ibid. D **84**, 049903 (2011)] [arXiv:0812.4208 [gr-qc]].
 - [36] N. Yunes, A. Buonanno, S. A. Hughes, Y. Pan, E. Barausse, M. C. Miller and W. Throwe, Phys. Rev. D **83**, 044044 (2011) [arXiv:1009.6013 [gr-qc]].
 - [37] Z. Zhang, N. Yunes, E. Berti, Phys. Rev. **D84**, 024029 (2011). [arXiv:1103.6041 [gr-qc]].
 - [38] H. Tagoshi, M. Shibata, T. Tanaka, M. Sasaki, Phys. Rev. **D54**, 1439-1459 (1996). [gr-qc/9603028].
 - [39] L. Blanchet, B. R. Iyer and B. Joguet, Phys. Rev. D **65**, 064005 (2002) [Erratum-ibid. D **71**, 129903 (2005)] [gr-qc/0105098].
 - [40] L. Blanchet, G. Faye, B. R. Iyer and B. Joguet, Phys. Rev. D **65**, 061501 (2002) [Erratum-ibid. D **71**, 129902 (2005)] [gr-qc/0105099].
 - [41] L. Blanchet, A. Buonanno and G. Faye, Phys. Rev. D **84**, 064041 (2011) [arXiv:1104.5659 [gr-qc]].
 - [42] R. Fujita, Prog. Theor. Phys. **127**, 583 (2012) [arXiv:1104.5615 [gr-qc]]. R. Fujita, to be published in Prog. Theor. Phys. **128**, (2012)
 - [43] R. Fujita, in preparation.
 - [44] H. Nakano, R. Fujita and N. Sago private communication.
 - [45] R. H. Boyer and R. W. Lindquist, J. Math. Phys. **8**, 265 (1967).
 - [46] E. A. Huerta and J. R. Gair, Phys. Rev. D **84**, 064023 (2011) [arXiv:1105.3567 [gr-qc]].
 - [47] E. A. Huerta, J. R. Gair and D. A. Brown, Phys. Rev. D **85**, 064023 (2012) [arXiv:1111.3243 [gr-qc]].
 - [48] J. M. Bardeen, W. H. Press, S. A. Teukolsky, Astrophys. J. **178**, 347 (1972).
 - [49] R. P. Geroch and J. Winicour, J. Math. Phys. **22**, 803 (1981).
 - [50] S. A. Teukolsky, Astrophys. J. **185**, 635-647 (1973).
W. H. Press and S. A. Teukolsky, Astrophys. J. **185**, 649 (1973).
S. A. Teukolsky and W. H. Press, Astrophys. J. **193**, 443 (1974).
 - [51] R. Fujita, H. Tagoshi, Prog. Theor. Phys. **112**, 415-450 (2004). [gr-qc/0410018];
R. Fujita, W. Hikida, H. Tagoshi, Prog. Theor. Phys. **121**, 843-874 (2009). [arXiv:0904.3810 [gr-qc]];

- [52] R. Fujita, H. Tagoshi, *Prog. Theor. Phys.* **113**, 1165-1182 (2005). [arXiv:0904.3818 [gr-qc]].
- [53] T. Damour, B. R. Iyer, B. S. Sathyaprakash, *Phys. Rev. D* **57**, 885-907 (1998). [gr-qc/9708034].
- [54] E. K. Porter and B. S. Sathyaprakash, *Phys. Rev. D* **71**, 024017 (2005) [gr-qc/0406038].
- [55] E. K. Porter, *Phys. Rev. D* **76**, 104002 (2007) [arXiv:0706.0114 [gr-qc]].
- [56] J. Carre and E. K. Porter, arXiv:1206.2509 [gr-qc].
- [57] T. Damour, B. R. Iyer and A. Nagar, *Phys. Rev. D* **79**, 064004 (2009) [arXiv:0811.2069 [gr-qc]].
- [58] R. Fujita and B. R. Iyer, *Phys. Rev. D* **82**, 044051 (2010) [arXiv:1005.2266 [gr-qc]].
- [59] Y. Pan, A. Buonanno, R. Fujita, E. Racine and H. Tagoshi, *Phys. Rev. D* **83**, 064003 (2011) [arXiv:1006.0431 [gr-qc]].
- [60] C. Cutler, E. Poisson, G. J. Sussman and L. S. Finn, *Phys. Rev. D* **47**, 1511 (1993).
- [61] M. Sasaki, H. Tagoshi, *Living Rev. Rel.* **6**, 6 (2003). [gr-qc/0306120].
- [62] H. Onozawa, *Phys. Rev. D* **55**, 3593 (1997) [gr-qc/9610048].
- [63] K. S. Thorne, *Astrophys. J.* **191**, 507 (1974).
- [64] J. L. Friedman, K. Uryu and M. Shibata, *Phys. Rev. D* **65**, 064035 (2002) [Erratum-ibid. *D* **70**, 129904 (2004)] [gr-qc/0108070].
- [65] E. Poisson, gr-qc/0410127.
- [66] H. Tagoshi, S. Mano and E. Takasugi, *Prog. Theor. Phys.* **98**, 829 (1997) [gr-qc/9711072].
- [67] K. Alvi, *Phys. Rev. D* **64**, 104020 (2001) [gr-qc/0107080].
- [68] E. Poisson, *Phys. Rev. D* **70**, 084044 (2004) [gr-qc/0407050].
- [69] S. A. Hughes, *Phys. Rev. D* **61**, 084004 (2000) [Erratum-ibid. *D* **63**, 049902 (2001)] [Erratum-ibid. *D* **65**, 069902 (2002)] [Erratum-ibid. *D* **67**, 089901 (2003)] [gr-qc/9910091].